

Nuclear/Radiological Terrorism: Emergency Department Management of Radiation Casualties

Jerrold T. Bushberg, PhD,*† Linda A. Kroger, MS, *† Marcia B. Hartman, MS,* Edwin M. Leidholdt, Jr, PhD,§† Kenneth L. Miller, MS,|| Robert Derlet, MD,‡ Cheryl Wraa, RN‡

†Department of Radiology, *Department of Environmental Health and Safety, ‡Department of Emergency Medicine, University of California Davis Health System, Sacramento, CA, §US Department of Veterans Affairs National Health Physics Program, Mare Island, CA, and ||Department of Radiology, Penn State Hershey Medical Center, Hershey, PA.

Adapted with permission from The Journal of Emergency Medicine, vol.32, No. 1, pp 71-85, 2007.

Reprint Address: Jerrold T. Bushberg, Ph.D., University of California Davis Health System, 2315 Stockton Blvd., FSSB 2500, Sacramento, CA 95817, jerrold.bushberg@ucdmc.ucdavis.edu

Abstract

Recent world events have increased concern that hospitals must be prepared for radiological emergencies. Emergency departments (EDs) must be ready to treat patients suffering from injuries in combination with radiation exposure or contamination with radioactive material. Every hospital should have a Radiological Emergency Medical Response Plan, tested through periodic drills, which will allow effective handling of contaminated and injured patients. Treatment of life-threatening or severe traumatic injuries must take priority over radiation related issues. The risk to ED staff from radioactive contamination is minimal if universal precautions are used. The likelihood of significant radiation exposure to staff under most circumstances is small. Educating medical staff on the magnitude of the radiological hazards allows them to promptly and

confidently provide the necessary patient care. Measures must be taken to prevent the “worried well” and uninjured people with radioactive contamination from overwhelming the ED.

Introduction

The concern that emergency department (ED) personnel will have to care for patients injured or contaminated by a terrorist act involving radioactive material has increased dramatically since the end of the cold war in the early 1990s. Sealed sources containing large amounts of radioactive material suitable for use by a terrorist are widely used in medicine and industry and many large sealed sources have been lost from governmental oversight, particularly in developing countries and states that were part of the former Soviet Union (1). This paper is intended to assist hospitals and emergency departments in preparing for nuclear/radiological terrorism and for accidents involving radiation injuries or radioactive contamination.

There are many possible terrorist radiation injury scenarios, including: (1) covert placement of a sealed radioactive source in a public location, (2) the use of a radiological dispersal device (RDD), (3) an attack on or sabotage of a nuclear facility, and (4) detonation of a nuclear weapon (atomic bomb), which is a remote but possible threat (1, 2). A sealed source may be covertly placed in a location that would expose many people until the presence of the source is discovered. While many people could be exposed, it is unlikely that any would be contaminated. An RDD is not a nuclear weapon; instead, it is a device designed to spread radioactive material for the purpose of terrorism. An RDD that uses a conventional explosive (e.g., dynamite or a plastic explosive) to spread the radioactive material is called a “dirty bomb.” While the initial explosion may kill or injure those closest to the bomb, the radioactive material that is dispersed will likely expose and contaminate survivors and emergency responders. Due to the limited dispersion by such an explosion, it is unlikely that the exposure or contamination of people outside the immediate blast area will have any clinical effect beyond the psychological impact from the fear of radiation and perhaps a slightly increased risk of cancer. To date, there has been no use of an RDD; however, the materials to produce an RDD have been intercepted by law enforcement (3).

An attack on or sabotage of a nuclear facility, such as a commercial irradiation facility or a nuclear power plant, could release large amounts of radioactive material. Detonation of a nuclear weapon (even one with a relatively low energy yield) in a populated area would result in extensive loss of life and wide-spread contamination. The use of a stolen or improvised nuclear device is the least likely scenario, because of the strict security controls over nuclear weapons and weapons-grade plutonium and uranium and the technical difficulty of constructing such a weapon, but the potential number of injured people is many times greater than from the other scenarios. In the case of a nuclear weapon detonation or an event resulting in the dispersion of used nuclear

reactor fuel, the radioactive contamination will consist of many radionuclides, including radioactive isotopes of iodine.

In all scenarios involving the release of radioactive material, radioactive contamination would be present at the site of the event. There may also be deposition of radioactive material by a downwind plume. The amount of contamination deposited by the plume would depend upon the nature of the event, the amount of radioactive material released, and atmospheric conditions such as wind speed.

There are several scenarios, other than those caused by terrorism, in which radiation emergencies may occur. Some examples include medical radiation therapy accidents; accidental overexposures from industrial irradiators; lost, stolen, or misused medical or industrial radioactive sources; accidents during the transportation of radioactive material; and nuclear reactor accidents. The Radiation Emergency Assistance Center Training Site (REAC/TS) recorded 428 major radiation accidents worldwide between 1944 and 2005, resulting in 126 radiation-related deaths (4). Though these accidents were infrequent, heightened awareness of the potential impact from terrorist activity has prompted many hospitals to reassess their preparedness for radiological emergencies.

Ionizing Radiation and Radioactive Material Definitions

Ionizing Radiation

Ionizing radiation, such as x and gamma rays as well as alpha and beta particles, has the ability to ionize matter, causing chemical changes that can modify DNA and kill cells. Ionizing radiation consists of energetic subatomic particles, such as alpha particles, beta particles, and neutrons, and ionizing electromagnetic radiation. Ionizing electromagnetic radiation is not to be confused with nonionizing electromagnetic radiation, such as visible light, infrared radiation, microwaves, and radio waves. Throughout this article, the term *radiation* will refer only to ionizing radiation. Radiation is a natural part of the environment. People are constantly exposed to radiation from their surroundings, (e.g., the earth itself, building materials, air, and water) as well as from cosmic rays. On average, persons in the United States receive about 300 millirem (mrem) of radiation per year from natural sources (5). People are also exposed to man-made radiation from medical imaging (e.g., radiology and nuclear medicine studies), medical therapy (e.g., cancer treatment), industry (e.g., soil moisture density gauges and nuclear power plants), and research.

Radioactive Material

Radioactive material consists of atoms with unstable nuclei. The nuclei of radioactive material spontaneously transform into more stable forms with the emission of ionizing

radiation. The amount of radioactivity is described by the quantity *activity*, defined as the number of nuclei undergoing transformation per unit time. The traditional unit of activity is the curie, defined in Table 1. Contamination of people usually involves microcurie (μCi ; one millionth of a curie) to millicurie (mCi ; one thousandth of a curie) quantities. Quantities of radioactive material range from trivial (e.g., μCi) amounts in typical biomedical research laboratories, to more than a billion curies in a commercial nuclear power reactor. Nuclear Medicine patients are injected with μCi to mCi quantities of short-lived radioactive material for routine diagnostic examinations. The amount of radioactivity continuously decreases with time, a phenomenon referred to as radioactive decay. The physical half-life is the time required for radioactive material to be reduced to half the initial amount by radioactive decay. After one half-life, half of the radioactive atoms remain. After a second half-life, a half of a half, or 25%, of the initial quantity of radioactive material remains, and so on. After ten half-lives, the radioactivity has been reduced by a factor of approximately 1000. The effective half-life combines the physical half-life with biological elimination (e.g., urination, defecation, exhalation, and sweating). Most of the radionuclides that are considered likely to be used in an RDD (see below) have physical half-lives of 5 years or more.

Table 1. Radiation Units

Quantity	Definition	Traditional Unit	SI Unit*	Unit Conversion
Activity	Amount of radioactivity; number of atoms undergoing nuclear transformations (nt) per time	curie (Ci) $1 \text{ Ci} = 37 \times 10^9 \text{ nt/s}$	becquerel (Bq) $1 \text{ Bq} = 1 \text{ nt/s}$	$1 \text{ Ci} = 37 \times 10^9 \text{ Bq}$
Absorbed dose	Energy imparted per mass	rad $1 \text{ rad} = 0.01 \text{ J/kg}$	gray (Gy) $1 \text{ Gy} = 1 \text{ J/kg}$	$100 \text{ rad} = 1 \text{ Gy}$
Effective dose	Sum of absorbed doses to organs, each multiplied by a tissue weighting factor	rem	sievert (Sv)	$100 \text{ rem} = 1 \text{ Sv}$
Exposure	Ionization in air per mass air	roentgen (R)	coulomb/kilogram (C/kg)	$1 \text{ R} = 2.58 \times 10^{-4} \text{ C/kg}$

Although international units (SI) are used by professionals in radiation protection, most instruments and many labels on sources of radioactive material still use traditional units.

For x and gamma radiation $1\text{R} \cong 1 \text{ rad} \cong 1 \text{ rem}$

Two common types of radioactive decay are alpha particle emission and beta particle emission. Alpha particles travel very short distances ($<0.1 \text{ mm}$) and therefore, are only

harmful when inhaled, swallowed or present in an open wound. Some beta particles can travel up to 10 meters or more in air and a centimeter or more in soft tissue. Thus, most beta emitting radioactive materials, if allowed to remain on the skin for a prolonged period of time, can cause skin injury. Beta emitting contaminants may also be harmful if deposited internally. Gamma radiation and x-rays are also emitted by many radioactive materials and travel many meters in air and many centimeters in living tissue. Layers of dense materials, such as lead, are commonly used to shield against gamma radiation and x-rays. Some of the radioactive materials considered likely to be used in an RDD are the beta emitters cesium-137 (Cs-137), cobalt-60 (Co-60), iridium-192 (Ir-192), strontium-90 (Sr-90) and iodine-131 (I-131) and the alpha emitters americium-241 (Am-241), californium-252 (Cf-252), and plutonium (Pu) (1, 6).

Quantities and Units for Describing Ionizing Radiation

The amount and rate of energy deposition in tissue by radiation are major determinants of biological effects. *Absorbed dose*, commonly referred to as *dose*, is defined as the amount of energy deposited per unit mass of matter. The traditional unit of dose is the rad (Table 1).

The biological effects of radiation also depend on the type of radiation. Some types of radiation, such as alpha particles and neutrons, are more biologically damaging per unit dose than x and gamma rays. The *effective dose* reflects the biological damage per unit dose (Table 1). The traditional unit of effective dose is the rem. For practical purposes when dealing with beta, gamma, and x radiation, 1 rad = 1 rem.

Contamination versus Exposure

The difference between radioactive contamination and radiation exposure is important to understand. Radioactive contamination is radioactive material that is in an unwanted location. In the case of an accident victim, it may be on the skin or clothes of the person or have entered the body by inhalation, ingestion, or through a wound. Contaminated patients require careful handling to effectively remove and limit the spread of contamination. Patients are not contaminated if they have only been exposed to ionizing radiation from a radioactive source or an x-ray machine. They do not pose any hazard to hospital personnel and radiation safety precautions are not needed.

Emergency Department and Hospital ProceduresProtection of Staff

Reducing Exposure

The three methods for reducing radiation exposure from a source external to the body to levels that are as low as reasonably achievable (ALARA) are time, distance and shielding. Radiation exposure rates decrease greatly (by the square of the distance) as you increase the distance from the source. For example, doubling the distance from the source decreases the radiation exposure rate to one quarter (Figure 1). The time spent in close proximity to the severely contaminated patient should be kept to a minimum while providing the necessary medical care. In general, exposure from contaminated patients will be minimal and, in most cases, lead aprons are not recommended.




Figure 1. The Inverse Square Law. Doubling the distance from a source of radiation reduces the exposure to one quarter. (Reproduced with permission from The Essential Physics of Medical Imaging, second edition, pg 757).

Protection from Contamination

Hospital staff are well versed in protecting themselves and their work areas from biological contamination through the use of “Universal Precautions.” The same precautions can be used effectively to protect personnel and the work area from contamination by radioactive materials. The only variation from normal universal precautions is a recommendation to wear two sets of gloves and to change the outer pair as needed to avoid cross contamination. It should be pointed out that in the case of a multi-hazard incident (e.g., radioactive materials and chemicals), higher levels of personal protective equipment may be necessary.

Detection and Measurement of Radiation

Hospitals with Nuclear Medicine or Radiation Oncology departments have radiation monitoring instruments. The ED should either have its own, or ready access to, such instruments that are properly maintained and calibrated. The ED training program should include instruction on instrument operation including the proper means of establishing the background levels. Background readings should be taken in an area of the ED remote from the areas that may be contaminated. The most commonly used instruments to detect the presence of radioactive material and radiation are the Geiger-Mueller (GM) survey meter (also known as a Geiger counter) and the ionization chamber survey meter (Figure 2).



Figure 2. A GM survey meter will detect small amounts of radioactive contamination and typically has the capability of distinguishing between alpha, beta and gamma radiation. It is useful in assessing where contamination is located and aids in the effectiveness of decontamination efforts.

The GM survey meter will detect small amounts of radioactive contamination and typically has the capability of distinguishing between alpha, beta, and gamma radiation. It is used to quickly determine if a person is contaminated and where the contamination is located. The detectors of GM survey meters to be used for locating radioactive contamination should have thin windows to permit the detection of alpha and beta particles, which are not as penetrating as gamma rays. GM survey meters may become inaccurate in very high radiation fields and therefore other instruments, such as ionization chamber survey meters, should be used to measure very high radiation fields. The radiation measurement of a GM survey meter is usually expressed as exposure per unit time (milliroentgen per hour; mR/hr) or counts per minute (CPM) (Figure 3).




Figure 3. GM survey meters (lower) are used for locating radioactive contamination but may become inaccurate in very high radiation fields. The radiation measurement of a GM survey meter is usually expressed as exposure per unit time (milliroentgen per hour; mR/hr) or counts per minute (CPM). Other instruments, such as ionization chamber survey meters (upper) are used to measure exposure rates from relatively low levels (mR/hr) to very high levels (R/hr) but are less sensitive than GM survey meters for detecting small amounts of contamination and are not recommended for this purpose.

Ionization chamber survey meters are used to measure exposure rates from low (mR/hr) to high (R/hr) levels of radiation. To determine the exposure a person actually receives, the measured exposure rate must be multiplied by the time that the person was exposed. For example, if staff were caring for a contaminated patient for 15 minutes and the ionization chamber reading was 5 mR/hr, the staff would be exposed to 1.25 mR, less than the radiation exposure received from cosmic radiation during a

cross-country airplane flight (approximately 2.5 mrem). The average natural radiation dose to people living in the US is 300 mrem per year. Examples of some typical radiation doses are listed in Table 2. Ionization chamber survey meters are less sensitive than GM survey meters for detecting small amounts of contamination and are not recommended for this purpose (Figure 3). Other specialized equipment, such as multi-channel analyzers (MCA) which allow radionuclide identification, might only be available if the hospital has health physics or medical physics staff.

Table 2. Typical Radiation Doses

Flight from Los Angeles to New York	2.5 mrem
Chest x-ray	5 mrem
Annual natural background in USA	300 mrem
Nuclear Medicine bone scan	400 mrem
Abdominal CT	800 mrem
Barium enema	900 mrem
Cardiac catheterization (skin dose)	45,000 mrem

A personal dosimeter is a device that measures the dose of radiation received by the person wearing it. Film badge dosimeters must be sent for analysis so the radiation dose received is not known for several days. Self-reading dosimeters, while not as accurate as film badges, allow the wearer to see the total radiation dose they have received at any time. Some of these are equipped with alarms that alert the wearer when a set point is exceeded. Ideally, both types of dosimeters should be used by staff working directly with contaminated patients.

Hospital Plan

The hospital should have a well thought out Radiological Emergency Medical Response Plan and should test it periodically by drills. This may be part of the emergency management plan required by the Joint Commission on Accreditation of Hospital Organizations (Joint Commission) (7). There are several excellent documents that can assist in plan development (2, 8, 9, 10, 11, 12, 13). The plan should ensure that the medical care of the patient takes priority over decontamination. Resuscitation and stabilization are the primary objectives with decontamination efforts being secondary. The plan should address contamination control for staff and facilities, including control and survey of materials and personnel entering and leaving the contaminated area (Table 3). Facility preparation depends on the notification time for patient arrival as well

as the number of patients expected. In situations involving other types of hazardous materials, such as chemicals, decontamination of the victims typically occurs prior to transportation to the ED. While this approach may be appropriate for some chemical and biological agents, it is unnecessarily restrictive for radiological contamination which poses very little risk of harm to the emergency responders. In addition, many casualties will likely self-refer to the nearest hospital and therefore not be decontaminated. The hospital plan should specifically address the issue and provide for admission of critical patients directly into the ED or related services (e.g., CT, surgery) without decontamination if a delay would be life threatening. Emergency medical technicians (EMTs) should attempt to decontaminate the victims in the field if the patients are stable, remembering that medical management is the first priority. Typically 90% of radioactive contamination is removed when the clothes are removed (14). What little remains will likely stay in place if the patient is wrapped in a sheet and transported.

Table 3. Facility Management of Radiological Emergencies

Activate Hospital Plan	1. Obtain radiation survey meters.
	2. Call for additional support; staff from Nuclear Medicine, Radiation Oncology, Radiation Safety (Health Physics).
	3. Establish triage area.
	4. Establish area for decontamination of uninjured persons.
Protect Staff from Contamination	1. Use universal precautions and double glove.
	2. Survey hands and clothing with radiation survey meter.
	3. Replace gloves and gowns if contaminated and between patients.
	4. Keep areas outside the treatment area free from contamination.
Plan to Control Contamination	1. Instruct staff to use universal precautions.
	2. Provide multiple containers for contaminated waste.
	3. Protect the floor with covering if time permits.

There should be a call list to notify the staff on duty, as well as to obtain additional staff and equipment that will be needed. The additional support for the ED could come from hospital staff in Nuclear Medicine, Radiation Oncology, Diagnostic Radiology, and Radiation Safety/ Health Physics. If radiation safety personnel are available, they will be a valuable asset in the management of the flow of people through the ED. They can perform surveys of the accident victims, assist in contamination control, and provide

radiation dose assessment (15). The plan should address where to obtain radiation survey meters. Diligent use of survey meters by staff (e.g., survey gloves and change as necessary, survey shoes when leaving, etc.) will help prevent the spread of contamination (Figure 2).

The plan should designate a triage area outside the ED. In this area, assessment can be made as to whether the person needs to be seen in the ED or not (2, 16). A mass casualty incident involving radioactive material is likely to generate large numbers of frightened people who do not require acute medical care (2, 14). The plan should provide for centers for assessment of minor injuries, decontamination, and counseling of these persons away from the ED so that the ED is not inundated. These centers should be staffed by physicians with radiological training, health physicists or other staff trained to use radiation survey instruments, and psychological counselors. The hospital should also plan to provide psychological support to patients and hospital staff.

Patient flow in the ED should be established in a manner that will control the spread of contamination (Figure 4) (9, 16, 17). Several methods to consider are: placing a floor covering on one side of the hallway from the ED entrance to the treatment room; designating a separate entrance for contaminated patients; and transferring the contaminated patient to a clean gurney before entering the ED. Ideally, the treatment room floor should be covered, but only if time allows for such an activity. Containers for waste should be set up to limit the spread of contamination. Compared to chemical and biological hazards, one advantage of radioactive contamination is the ease with which it can be detected. Frequent use of the GM survey meter can alert personnel of the need to change their gloves or clothing when they become contaminated and to identify when contamination is being spread so that cleanup and extra precautions can be implemented. Contamination in the ED may occur and this should not preclude the continued use of the treatment areas or equipment during the event.

The plan should identify a laboratory that can assay samples collected to assess external and internal contamination (Table 4). The laboratory should be capable of identifying the radionuclide(s) involved and estimating the amounts of radioactive material in the samples, particularly in regards to possible internal contamination (discussed below under "Assessment and Therapy for Internal Contamination"). A gamma ray spectroscopy system with an MCA is particularly useful for identifying radionuclides. A medical center with full time radiation safety staff will likely have this capability and a Nuclear Medicine department can develop this capability. Arrangements should be made in advance so that the laboratory can calibrate its equipment for the radionuclides likely to be encountered and provide staff training.




Figure 4. Patient flow through the ED. Familiarization of the ED staff to the hospitals Emergency Management Plan through the use of periodic drills will maximize their effectiveness and allow for optimal care of the patients being seen.

Table 4. Specimens to be Collected for Medical Assessment of a Patient Exposed to Radiation

Specimen/Type of Analysis	Reason for Obtaining	Mechanism for Obtaining
<i>In All Cases of Radiation Injury</i>		
Blood: CBC with differential lymphocyte count, repeated every 6 hrs for 48 hrs if possibility of total body irradiation.	To establish a baseline and change over time. A significant decrease in lymphocyte count is an early predictor of radiation dose. See Table 8.	Veni-puncture in uncontaminated area into purple top tube containing EDTA; cover puncture site.
Blood: Chromosomal analysis	Chromosomal analyses are the most accurate way of estimating the radiation dose. Specialized labs are required and results take several days. See Table 8.	Veni-puncture in uncontaminated area into lithium heparin tube; if that is not available, use an EDTA tube. Cover puncture site.
Urine: Routine urinalysis	Determine normal kidney function and baseline for urinary constituents; especially important if internal contamination is suspected	Avoid contamination when collecting sample. Label sample with date and time of collection.
<i>When External Contamination is Suspected</i>		
Nose, ear, mouth: Swab body orifices. Analyze with GM probe, gamma or liquid scintillation counter, or multichannel analyzer if available.	Assess the possibility of internal contamination and identify the radionuclide.	Use separate saline or water moistened swabs to wipe the inside of each nostril, ear, and mouth. Label and bag separately.
Wounds: Samples from dressings or swabs of wounds. Analyze with GM probe, gamma or liquid scintillation counter, or multichannel analyzer if available.	Assess if wounds are contaminated and identify the radionuclide.	Save dressings as they are changed. Use swabs to sample the secretions from wounds. If foreign objects or debris are removed from the wound or skin, transfer it to lead containers (pigs).

<i>When Internal Contamination is Suspected</i>		
Urine Bioassay: 24 hr specimen; repeat for 4 days	Body excreta may contain radionuclides if internal contamination has occurred.	Use standard specimen containers.
Feces Bioassay: 24 hr specimen; repeat for 4 days		

Adapted from Radiation Emergency Assistance Center/Training Site (REAC/TS) (4). *Hospital emergency care of the radiation accident patient*. 2002. <http://www.orau.gov/reacts/emergency.htm>

Patient Assessment and Management of Radioactive Material Contamination and Radiation Injuries

Patient Assessment

Questions about the circumstances surrounding the incident as well as a GM survey of the patient will provide valuable patient management information and will further assist in predicting the extent of radiation injury (9, 13, 18). Although contamination surveys are secondary to patient stabilization, they should be conducted as soon as possible. In the case of an RDD, very high dose rates could exist if embedded shrapnel from the RDD is intensely radioactive, which is easily determined with radiation survey equipment. This type of shrapnel should be transferred to lead containers (readily available from Nuclear Medicine departments and commonly referred to as “pigs”) and placed distant from staff. Specimens collected for medical assessment can yield valuable information for treatment planning (Table 4). Samples collected to assess contamination should be placed in containers and labeled with the patient’s name, date, time of collection and sample location. ED staff should be aware that elevated radiation measurements from a patient may be caused by radioactive material from a recent nuclear medicine or radiation oncology procedure instead of radioactive contamination from an event. Nausea and vomiting are signs of exposure to a high radiation dose (see Acute Radiation Syndrome section below), but such a dose from terrorist activities is unlikely. Consequently, if such symptoms are present, they may be psychological in nature.

Decontamination

Decontamination should not delay or impede stabilization of the patient (Table 5). Removal of clothing generally reduces contamination on the patient by 90% (14). Usual washing methods are effective for removal of radioactive contamination. Non-contaminated wounds should be protected with waterproof dressings to minimize uptake of radioactive material during decontamination efforts. Contaminated waste water need not be contained if it will unduly

complicate the treatment of the patient or if it is otherwise determined to be impractical. Decontamination efforts should not cause the skin to become abraded. Openings in the skin

Table 5. Step - By - Step Patient Decontamination

1. Do not delay surgery or other necessary medical procedures or exams; residual contamination can be controlled.
2. Carefully remove and bag patient's clothing and personal belongings (typically removes 90% of contamination).
3. Survey patient and, if practical, collect samples.
4. Handle foreign objects with care until proven non-radioactive with a radiation survey meter.
5. Priorities <ul style="list-style-type: none"> A. Decontaminate wounds first then intact skin. B. Start with the highest levels of contamination. C. Protect non-contaminated wounds with water-proof dressings. D. Irrigate and gently scrub contaminated wounds with a surgical sponge. E. Extend wound debridement for removal of contamination only in extreme cases and upon expert advice. F. Change dressings frequently.
6. Decontaminate thermal burns by gently rinsing. Aggressive washing may increase the severity of injury. Additional contamination will be removed when dressings are changed.
7. Decontaminate intact skin and hair by washing with soap and water.
8. Remove stubborn contamination on hair by cutting with scissors or electric clippers.
9. Promote sweating for residual skin contamination.
10. Avoid overly aggressive decontamination.
11. Cease decontamination of skin and wounds: <ul style="list-style-type: none"> A. When the area is less than twice background, or B. When there is no significant reduction between decontamination efforts and before the skin becomes abraded.
12. Change outer gloves frequently to minimize spread of contamination.
13. Use survey meter to monitor progress of decontamination.

allow increased absorption of radioactive material. If an area of contaminated skin persists, cover the area with gauze and a glove or plastic to promote sweating which can remove radioactive material from pores. Remove contaminated hair, if necessary, using scissors or electric clippers. To avoid cutting the skin, do not shave.

To decontaminate wounds, irrigate with tepid water and gently wash with soap and a surgical sponge or gauze pad. Normal wound debridement can be performed. Excision around wounds solely to remove contamination should only be performed in extreme cases and upon the advice of radiological emergency medical experts. Contaminated thermal burns can be gently rinsed, but should not be scrubbed, to ensure that there will be no further damage to the skin. Often radioactive material will exude from wounds into gauze dressings, so frequent changing of dressings may aid wound decontamination. The dressing also serves to keep the contamination from spreading. Cease decontamination of the skin and wounds when the area is less than twice the background reading on the GM survey meter or if there is no significant reduction between washes.

Under no circumstances should emergency surgery or other necessary medical procedures be delayed because of contaminated skin or wounds. Staff will be protected from becoming contaminated by using universal precautions. Sheets and dressings will keep contamination in place. When the patient is ready to be moved from the ED to other areas of the hospital, the patient can be transferred to a clean gurney. The clean gurney can be brought into the contaminated treatment room by rolling it over clean paper or other clean material placed on the floor. After the patient is transferred onto the gurney, wrap a clean sheet around the patient and he or she can be transported throughout the hospital without spreading radioactive contamination.

Assessment and Therapy for Internal Contamination

Deposition of radioactive materials in the body (i.e., internal contamination), is a time-dependent, physiological phenomenon related to both the physical and chemical natures of the contaminant. Internal contamination occurs mainly by inhalation, ingestion, or absorption through wounds, although percutaneous absorption occurs with some radioactive materials. The fraction of the radioactive material that is absorbed into the body depends upon the radionuclide, chemical and physical form, and route into the body. For example, soluble plutonium is readily absorbed into the body through the lungs or a wound, but only very slightly if ingested. Furthermore, the amount of radioactive material that poses a significant threat if taken into the body varies greatly with radionuclide and chemical form. For example, 100 μCi of Cs-137 chloride in the body would not pose a significant threat, but less than 1 μCi of soluble plutonium would. In many cases, early treatment is necessary if it is to have much effect. Thus, time can be critical and treatment decisions may need to be made based on preliminary information. If internal contamination is suspected, plan for 24 hour urine and fecal

collections (Table 4) but the decision to treat suspected internal contamination should not be delayed until these samples are collected or analyzed (19).

If an MCA is available, it will be a valuable tool to expedite the determination of which decorporation therapy is appropriate based on the radionuclide(s) present. Also, if inhalation of radioactive material is suspected, immediately take nasal swabs. A separate swab should be obtained from each nostril and placed in its own container. Radioactivity on both nasal swabs should be considered evidence of inhalation of radioactivity, but lack of radioactivity on the swabs should not lead to the presumption that such material was not inhaled. Several methods of preventing incorporation (e.g., catharsis, gastric lavage) might be applicable, depending on the type of radioactive material present.

NCRP Report No. 65, *Management of Persons Accidentally Contaminated with Radionuclides* (17), the *Guidebook for the Treatment of Accidental Internal Radionuclide Contamination of Workers* (20), and *Guidance for Industry, Internal Radioactive Contamination – Development of Decorporation Agents* (21) provide clinical guidance on the management of internal contamination. The NCRP is currently working on an update to Report No. 65 which will contain additional information. REAC/TS has medical experts on call 24 hours a day to provide assistance with issues such as decorporation and treatment of exposed individuals (4). Some of the medications or preparations used in decorporation (trisodium calcium diethylenetriaminepentaacetate [Ca-DTPA], Zn-DTPA, Prussian Blue, etc.) might not be available locally and should be stocked (or a source identified in the community) as part of the hospital's Radiological Emergency Medical Response Plan (Table 6). The US Food and Drug Administration has stated that Prussian Blue is safe and effective in treating people exposed to cesium-137 and that Ca-DTPA and Zn-DTPA are safe and effective for treatment of internally deposited plutonium, americium, and curium (21). If it is suspected that the patient may have been exposed to radioiodine, potassium iodide (KI) should be administered to help protect the thyroid gland (Table 7) (17, 20, 22, 23). It should be noted that, with the exception of KI which is very effective if used early after exposure to radioiodines, most decorporation agents will only reduce the effective dose from radiation to the patient by a factor of 3 or less.

Table 6. Substances Suitable for Decorporation Treatment

Medication	Trade Name	Elements	Dosage Schedule*	Relative Effectiveness with Time	Principle of Action	Side Effects	Contraindications
DTPA (trisodium diethylene triaminepentaacetate, (Ca or Zn) (ref 21)	N/A	Am, Cf, Cm, Pu	Adults: 1 g/d by slow IV push or infusion in 250 ml 5% glucose. Inhalation in a nebulizer: 1 g/d Pediatrics (less than 12 years): 14 mg/kg/d by slow IV push or infusion in 250 ml 5% glucose Ca-DTPA for initial treatment (24 hr) followed by Zn-DTPA for protracted therapy.	Should be started as soon as possible. Efficacy greatest within 6 hr of exposure.	Chelation	Monitor blood pressure during infusion. Monitor for kidney function, liver enzymes, and signs of intestinal damage. Can be used for wound cleaning. May need trace metal supplements for prolonged therapy.	Ca-DTPA contraindicated for patients with kidney, intestinal or hematopoietic disorders, pregnant women, and minors.
KI (potassium iodide)	Thyroblock, Iostat and others	I	130 mg/day oral†	50% effective at 6 hr. Only 7% effective if given at 24 hr.	Blocking	Angioedema if sensitive to iodine; Treat w/ perchlorate, 1 x 200 mg, then 100 mg every 5 hr.	Caution with goiter or hyperthyroidisms; Do not treat the elderly.
Prussian Blue (ferric hexacyanoferrate) (ref 21)	Radiogardase	Cs	Adults and adolescents: 3 grams orally three times daily. Pediatrics (2-12 years): 1 gram orally three times daily.	Most effective when given promptly but continues to be effective in preventing recycling of radionuclide in the intestine.	Adsorption	Constipation. May affect absorption of other medications including tetracycline.	Only effective if GI motility is intact.

Radiation Casualties in the ED

<p>A) Aluminum phosphates</p> <p>B) Alginates</p> <p>C) NH₄Cl (ammonium chloride) plus Calcium gluconate</p>	<p>A) Aluminum containing antacids</p> <p>B) Gaviscon</p>	Sr	<p>A) 100 ml gel (13 g) oral</p> <p>B) 10 g oral then 4 g per day</p> <p>C) 1-2 g NH₄Cl every 6 hrs oral plus 500 mg Ca gluconate in 500 ml 5% glucose in water over 4 hr iv.</p>	<p>A & B) Most effective if given immediately.</p> <p>C) Still effective up to 2 wks after exposure.</p>	<p>A & B) Adsorption</p> <p>C) De-mineralization</p>	<p>A&B) Constipation. Possible interactions with other medications including tetracycline.</p> <p>C) Gastric irritation.</p>	<p>A&B) Diabetics: May contain glucose.</p> <p>C) Ca gluconate should not be given to patients with heart conditions.</p>
NaHCO ₃ (sodium bicarbonate)	N/A	U	4 g oral initially; 2 g every 4 hr until urine pH of 8-9 is maintained.		Complexation	None	
Penicillamine	Cuprimine, Depen	Heavy Metals including Cu, Fe, Hg, Pb, Au, Po	250-1500 mg every 8 hr oral		Chelation	Maculopapular or erythematous rash, marrow suppression, kidney function, immune-related diseases.	

* Consult REAC/TS (4) or the Physicians Desk Reference (PDR) for information on pediatric dosages not provided.

† See Table 7 for additional information.

Table 7. Thyroid Radioiodine Predicted Dose Threshold for Administration of KI for Different Risk Groups (22)

	Predicted Thyroid Dose (rad)	KI dose (mg)	Number of 130 mg tablets*	Number of 65 mg tablets*
Adults over 40 yrs	≥500	130	1	2
Adults 18-40 yrs	≥10	130	1	2
Pregnant or lactating women	≥5	130	1	2
Adolescents 12-18 yrs†	≥5	65	½	1
Children 3-12 yrs	≥5	65	½	1
1 mo-3 years	≥5	32	¼	½
birth - 1 month	≥5	16	1/8	¼

*Take tablet or fraction of tablet orally once a day. Crush for small children. Saturated Solution of Potassium Iodide (SSKI) can be used in place of tablets. SSKI contains 1 gm of KI per ml of solution. 130 mg of SSKI = 0.13 ml, or approximately 3 drops. Dilute in juice.

† adolescents approaching adult size (> 70 kg) should receive the full adult dose (130 mg)

Acute Radiation Syndrome

Acute radiation syndrome (ARS) (i.e., radiation sickness) is rare but may present as an acute illness in patients following exposure to very large doses (greater than 100 rem) of radiation. ARS is extremely unlikely to occur in staff treating patients who are contaminated with radioactive material. ARS follows a roughly predictable course over a period of time ranging from a few hours to several weeks (Table 8) and is described in many references (9, 14, 24). The severity of the syndrome increases with radiation dose, amount of the body exposed (whole body vs. partial body exposure), and the penetrating ability of the radiation. The severity is also affected by factors such as age, genetics, and medical conditions. If it is believed that the victim may have received a high radiation dose (e.g., greater than 50 rem), a complete blood cell count with absolute lymphocyte count should be taken initially and about every six hours thereafter (Table 4). The higher the radiation dose to the patient, the more severe is the drop in the concentration of circulating lymphocytes. The concentration of lymphocytes in circulation can be altered by trauma, which can complicate the use of blood cell concentrations as an indicator for radiation exposure. If chromosomal analyses are ordered, lithium heparin tubes should be used to collect the blood samples; lithium heparin with gel is not acceptable. If lithium heparin tubes are not available, use EDTA tubes. Chromosomal analyses of lymphocytes are performed by specialized laboratories and typically require days for results to be reported. REAC/TS can assist with locating the appropriate facilities (4). Results of these tests may be used to

estimate the radiation dose and the severity of radiation injury.

Table 8. Phases of Acute Radiation Syndrome

Phase of Syndrome	Feature	Effects of whole-body irradiation, from external radiation or internal absorption, by dose range in rad (1 rad = 1cGy; 100 rad = 1Gy)					
		0-100	100-200	200-600	600-800	800-3000	> 3000
Prodromal	Nausea, vomiting	None	5-50%	50-100%	75-100%	90-100%	100%
	Time of onset		3-6 hr	2-4 hr	1-2 hr	< 1 hr	Minutes
	Duration		< 24 hr	< 24 hr	< 48 hr	< 48 hr	N/A
	Lymphocyte count	Unaffected	Minimally decreased	< 1000 at 24 hr	< 500 at 24 hr	Decreases within hours	Decreases within hours
	CNS function	No impairment	No impairment	Routine task performance Cognitive impairment for 6-20 hr	Simple, routine task performance Cognitive impairment for > 24 hr	Rapid incapacitation May have a lucid interval of several hours	
Latent	No symptoms	> 2 wk	7-15 d	0-7 d	0-2 d	None	
Manifest illness	Signs, symptoms	None	Moderate leukopenia	Severe leukopenia, purpura, hemorrhage, pneumonia, hair loss after 300 rad		Diarrhea, fever, electrolyte disturbance	Convulsions, ataxia, tremor, lethargy
	Time of onset		> 2 wk	2 d - 2 wk		1-3 d	
	Critical period		None	4-6 wk; greatest potential for effective medical intervention		2-14 d	1-48 hr
	Organ system	None		Hematopoietic, respiratory (mucosal) systems		GI tract, Mucosal systems	CNS
	Hospitalization Duration	0%	< 5% 45-60 d	90% 60-90 d	100% 90+ d	100% Weeks to months	100% Days to weeks
	Mortality	None	Minimal	Low with aggressive therapy	High	Very high; significant neurological symptoms indicate lethal dose	

From Military Medical Operations Armed Forces Radiobiology Research Institute. *Medical Management of Radiological Casualties Handbook*. Second Edition. April 2003. (14) <http://www.afri.usuhs.mil>

The signs and symptoms that develop in ARS occur in four distinct phases: prodromal (initial), latent period, manifest illness stage, and recovery or death (Table 8). During the prodromal phase, symptoms, including loss of appetite, nausea, vomiting, fatigue, and diarrhea, should be treated in a routine clinical manner. The severity and time of onset of symptoms should be recorded. The severity and time of onset of symptoms should be recorded. Higher radiation doses increase the severity and the promptness of symptom onset. The time to vomiting, lymphocyte depletion kinetics and presence of chromosome dicentrics have been found to be predictors in estimating external radiation exposure and helpful to use to triage patients arriving at the ED (24).

Following the prodromal phase, there is a latent period in which the patient appears to recover. The duration of the latent phase decreases as the radiation dose increases. For doses in the range of 100 to 800 rem to most of the body, the latent period ranges from weeks to a few days and the manifest illness stage begins in less than 2 weeks and is mainly caused by damage to the hematopoietic system, resulting in decreased blood concentrations of leukocytes and platelets, which cause an increased risk of infection and bleeding. For healthy people without medical intervention, a dose of about 200 rem will result in less than 10% mortality while a dose of 400 rem will result in approximately 50% mortality (12).

For doses above about 800 rem, the latent period is on the order of one to three days and the primary effects are due to damage to the lining of the intestines, with diarrhea, fever, sepsis, and electrolyte disturbances. Survival, even with medical treatment, is highly unlikely for doses above about 1,000 rem. For even higher doses, exceeding about 3,000 rem, there is little or no latent period. The patient soon becomes confused, with severe diarrhea and hypotension, and death occurs in a day or two.

Treatment of ARS should focus on prevention of infection. Antibiotics for neutropenia and fever should be guided by the recommendations of the Infectious Diseases Society of America (25). Routine gut prophylaxis should not be administered (24). Hematopoietic growth factors should be given within the first 24-48 hours and then daily. Patients with severe ARS will require hospitalization. Several books and papers provide guidance on the management of the ARS (2, 9, 14, 24). REAC/TS and the Medical Radiobiology Assistance Team of the Armed Forces Radiobiology Research Institute (AFRRI) can provide guidance 24 hours a day on patient assessment and management of ARS (Table 9).

Table 9. Resources

Organizations

Radiation Emergency Assistance Center/ Training Site	(865) 576-1005	www.orau.gov/reacts
Centers for Disease Control (CDC)	(888) 246-2675	www.bt.cdc.gov/radiation/
Medical Radiobiology Advisory Team (MRAT) of the Armed Forces Radiobiology Research Institute (AFRRI)	(301) 295-0530	www.afri.usuhs.mil

Books

Medical Management of Radiation Accidents, Gusev, Guskova and Mettler, CRC Press, 2001.
Medical Effects of Ionizing Radiation, Mettler and Upton, Saunders, 1995.
The Medical Basis for Radiation-Accident Preparedness, REAC/TS Conference, 2002.
National Council on Radiation Protection and Measurements Report Nos. 65 and 138.
Distribution and Administration of Potassium Iodide in the Event of a Nuclear Incident, National Academy of Sciences (http://books.nap.edu/catalog/10868.html). 2004.

Booklets/Articles

Disaster Preparedness for Radiology Professionals, Response to Radiological Terrorism (http://www.acr.org)
Guidebook for the Treatment of Accidental Internal Radionuclide Contamination of Workers. Radiation Dosimetry, 41:1-49, 1992. Department of Health and Human Services, Food and Drug Administration, Guidance for Industry Internal Radioactive Contamination – Development of Decorporation Agents, March 2006 (http://www.fda.gov/cder/guidance/index.htm)
Department of Homeland Security Working Group on Radiological Dispersal Device Preparedness, (http://www1.va.gov/emshg/docs/Radiologic_Medical_Countermeasures_051403.pdf), 2003.
Interim Guidelines for Hospital Response from a Radiological Incident, (http://www.bt.cdc.gov/radiation/pdf/MassCasualtiesGuidelines.pdf), 2003.
“Major Radiation Exposure - What to Expect and How to Respond,” Mettler and Voelz, New England Journal of Medicine, 346:1554-61, 2002.
“Medical management of the acute radiation syndrome: recommendations of the strategic national stockpile radiation working group” Waselenko JK, MacVittie TJ, et al. Annals of Internal Medicine, 140: 1037-1051, 2004.
Medical Management of Radiological Casualties Handbook, (http://www.afri.usuhs.mil), 2003.
OSHA Best Practices for Hospital-Based First Receivers of Victims from Mass Casualty Incidents Involving the Release of Hazardous Substances, (http://www.osha.gov/dts/osta/bestpractices/firstreceivers_hospital.html), 2004.

Training Program

“Emergency Department Management of Radiation Casualties” Bushberg, JT, et al, (http://hps.org/hsc/documents/emergency.ppt), 2004.

REAC/TS and AFRRRI have developed a software application, the Biodosimetry Assessment Tool (BAT), which provides a database to record, integrate, and archive diagnostic information (clinical signs and symptoms, physical dosimetry, etc.) that is relevant to the management of radiation casualties. The BAT program is available on the AFRRRI website. Additionally, the AFRRRI website has many resources and worksheets that can be used to document this information.

Patients who have suffered trauma combined with an acute high level radiation exposure will have increased morbidity as compared to patients who have received the same dose of radiation without trauma. If a patient has received an acute radiation dose greater than 100 rem, efforts must be made to close wounds, cover burns, reduce fractures, and perform surgical stabilizing and definitive treatments within the first 48 hours after injury. After 48 hours, surgical interventions should be delayed until hematopoietic recovery has occurred.

Local Radiation Injury

Radiation exposure to parts of the body can cause localized effects if the dose is sufficiently high, typically greater than 200 rem. The patient may not be aware that he or she was exposed. Such a patient may have localized burn-like skin injuries without a history of heat exposure. Epilation of the exposed area may occur. These symptoms do not appear immediately but rather days after exposure. A tendency to bleed, nausea and vomiting or other symptoms of the ARS may be present. If a patient presents with burns immediately after a terrorist event, such as a dirty bomb, it is more likely that the burns are thermal burns, not radiation burns.

Other Health Effects of Radiation

Studies of animals and epidemiologic studies of humans, particularly the survivors of the atomic bombings in Japan, have shown that radiation is a weak carcinogen. The International Commission on Radiological Protection has estimated the lifetime risk of fatal cancer, for working age adults exposed to low doses of radiation, to be about 5×10^{-4} deaths per rem (26). For doses likely to be received by staff in the ED (<1 rem), the increased risk of cancer is minimal.

Radiation-induced hereditary (genetic) effects have been demonstrated in animals, but direct evidence of radiation-induced hereditary effects has not been observed in any of the studies of humans exposed to radiation, even after high doses of radiation (27). For doses likely to be received by staff, the risk is minimal. Using a theoretical model, the increased risk of genetic effects to children of young emergency responders would be ~0.02% from 5 rem and ~0.2% from 50 rem (27). The natural incidence of genetic disease and malformations at 1 to 2 years of age is 6-10% (28).

Studies of animals and humans have shown that prenatal radiation exposure can cause teratogenic effects (27, 29, 30, 31). However, there is no evidence of an increased incidence of such effects for doses less than 10 rem to the embryo or fetus. Also, some epidemiological studies of humans indicate that radiation-exposure to the embryo or fetus may increase the incidence of post-natal cancer (29). Although it is quite unlikely that someone providing care for a radioactively contaminated patient would receive a large dose, as a precaution, the CDC recommends excluding pregnant staff from such cases whenever possible (8). Fetal doses would have to be greater than 10 rem before abortions were considered and, then, only on advice from a physician with expertise on radiation and pregnancy (32).

Radiation Dose Limits For Emergency Workers

There are several recommendations for radiation dose limits to emergency workers performing life saving actions. While the US Nuclear Regulatory Commission (NRC) does not have a limit for emergency life saving actions (i.e. immediate response to a life or death situation), the National Council on Radiation Protection and Measurements (5) and the International Council on Radiation Protection (26) recommend that doses can approach or exceed 50 rem. While this dose is 10 times higher than annual occupational dose limit, it represents a modest increase in cancer risk compared to the large benefit to the recipients of the life saving measures. Emergency workers should perform these lifesaving activities voluntarily and with prior training regarding the risks of exposure. It is unlikely that the dose to ED staff would approach these levels. The doses ED staff received after the Chernobyl accident were less than 1 rem (12).

Facility Recovery

If the efforts made during facility preparation were successful, facility recovery should be relatively easy. During facility recovery, employee exposures should be kept ALARA. If an in-house radiation safety staff is available, they will supervise decontamination efforts. Waste from the ED and triage area should be taken to a designated holding location until it can be surveyed for radioactive material prior to disposal. Some facilities have portal radiation monitors to survey hospital trash. They provide a quick method that can replace hand surveying of each bag of waste. A radiation survey of the facility will identify any surfaces that require decontamination. Normal cleaning methods are typically very effective. Facilities should be decontaminated to the extent possible. Gloves, shoe covers, and coveralls should be worn by people decontaminating the area. For some contaminated items, replacement may be more cost effective and practical than decontamination. The decontamination goal for surfaces is to have less than twice the normal background reading. Areas that cannot be decontaminated to this level should be referred to health physics or radiation safety specialists. There are companies that can perform decontamination services.

Summary

In a radiological emergency, medical stabilization and treatment of the patient take priority over decontamination efforts. Radiation exposure and contamination are not likely to be significant hazards to staff. Staff can protect themselves from radioactive contamination by using universal precautions while treating these patients. As opposed to patients who arrive contaminated with chemical or biological agents, radioactive contamination is easy to detect.

Every hospital should have a Radiological Emergency Medical Response Plan that will allow effective handling of contaminated and injured patients. The Joint Commission requires organizations to have an emergency management plan (7). There are many resources available including books, journal articles, and internet sites that will be useful in preparing a hospital emergency plan (Table 9). Testing the plan through periodic drills (e.g., annually) and the utilization of an effective training program (Table 9) minimizes the potential for apprehension and panic should activation of the plan ever be needed. The plan should address patient assessment and management of radioactive contamination and radiation injuries. Training should emphasize that resuscitation and stabilization are the most important aspects of treating the radiation accident victim(s). Preparations should ensure that adequate supplies and radiation survey instruments are available. Non-ED staff that can assist in a radiological emergency should be identified and trained in advance. Staff from Nuclear Medicine, Radiation Oncology, and Radiation Safety have expertise in radiation protection practices and the use of survey meters. Measures should be taken to prevent the “worried well” and uninjured people with radioactive contamination from overwhelming the ED and hindering patient care.

Most victims of a mass casualty event involving radioactive material will arrive at the ED with little contamination. ED staff are likely to receive far less than the annual occupational dose limit of 5 rem. The long term risks from radiation exposures of less than 5 rem are very low. Even in an extreme case involving radiation casualties, the Chernobyl nuclear reactor accident in 1986, medical personnel working on the victims received less than 1 rem (12). In the case of an RDD, very high dose rates could exist if embedded shrapnel from the RDD was intensely radioactive. This unlikely possibility would be easy to determine with radiation survey instruments.

Most victims of radiation accidents will show no signs or symptoms of radiation exposure due to their small exposures. In the rare instance when victims have received large radiation doses, early signs and symptoms and their intensities will be an indication of the severity of the radiation injury. The first 24 hours after an incident will be the most challenging for those involved in the emergency response. After the first day, there will likely be many additional resources arriving from state and federal agencies. Understanding basic radiation protection principles will aid the care providers to effectively and efficiently managing the victims of radiation accidents.

Acknowledgments: The authors thank Fred A. Mettler, Jr., M.D., and William E. Dickerson, M.D. for their reviews of this manuscript.

References

1. Ferguson, CD, Potter WC. In: Donohoe LS ed. *The four faces of nuclear terrorism*. Monterey: Monterey Institute of International Studies; 2004:10-12.
2. National Council on Radiation Protection and Measurements Report No. 138. *Management of terrorist events involving radioactive materials*. Bethesda: NCRP Publications; 2001.
3. Center for Nonproliferation Studies at the Monterey Institute of International Studies. *Radiological Terrorist Tutorial: History of Radiological Terrorism*. Available at: http://www.nti.org/h_learnmore/radtutorial/chapter03_01.html Accessed: May 22, 2006.
4. Oak Ridge Institute for Science and Education. *Radiation Emergency Assistance Center/ Training Site (REAC/TS)*. Available at: www.ornl.gov/reacts/ Accessed: May 22, 2006.
5. National Council on Radiation Protection and Measurements Report No. 116. *Limitation of exposure to ionizing radiation*. Bethesda: NCRP Publications; 1993.
6. Centers for Disease Control and Prevention, Department of Health and Human Services. *Radioactive Isotopes*. Available at <http://www.bt.cdc.gov/radiation/isotopes/> Accessed: May 22, 2006.
7. Joint Commission on Accreditation of Healthcare Organizations. *The 2007 Joint Commission Accreditation Manual for Hospitals*. Oakbrook Terrace: Joint Commission on Accreditation of Healthcare Organizations; 2007.
8. Centers for Disease Control and Prevention, Department of Health and Human Services. *Interim guidelines for hospital response to mass casualties from a radiological incident*. Available at: www.bt.cdc.gov/radiation/pdf/masscasualtiesguidelines.pdf Accessed: May 22, 2006.
9. Mettler FA. In: Gusev IA, Guskova AK, Mettler FA, eds. *Medical management of radiation accidents* (2nd Ed). Boca Raton: CRC Press; 2001:425-35.
10. Bushberg JT, Miller, KL. In: Brodsky A, Johnson Jr RH, Goans RE, eds. *Public protection from nuclear, chemical and biological terrorism*. Madison, WI: Medical Physics Publishing; 2004:445-460.

11. *Occupational Safety and Health Administration. OSHA best practices for hospital-based first receivers of victims from mass casualty incidents involving the release of hazardous substances. January 2005. Available at: http://www.osha.gov/dts/osta/bestpractices/firstreceivers_hospital.html Accessed: May 22, 2006.*
12. *Mettler FA, Voelz GL. Major radiation exposure - what to expect and how to respond. N Eng J Med. 2002;346:1554-61.*
13. *International Atomic Energy Agency and World Health Organization. Generic procedures for medical response during a nuclear or radiological emergency. April 2005. Available at: http://www-pub.iaea.org/MTCD/publications/PDF/EPR-MEDICAL-2005_web.pdf Accessed: May 22, 2006.*
14. *Armed Forces Radiobiology Research Institute. Medical management of radiological casualties. Bethesda, MD: Armed Forces Radiobiology Research Institute, Second Edition; 2003. Available at: www.afrrri.usuhs.mil Accessed: May 22, 2006.*
15. *Toohy RE. Role of the health physicist in dose assessment. In: Ricks RC, Berger ME, O'Hara, Jr. FM, eds. Medical basis for radiation-accident preparedness. New York: The Parthenon Publishing Group, Inc.; 2002:33-43.*
16. *Miller KL. The regional approach to management of radiation accident victims. In: Vince MA, ed. Coping with radiation accidents. Ravenna: Envirotox Management, Inc.; 1990:121-30.*
17. *National Council on Radiation Protection and Measurements Report No. 65. Management of persons accidentally contaminated with radionuclides. Bethesda: NCRP Publications; 1979.*
18. *Miller KL, DeMuth WE. Handling radiation emergencies: no need to fear. J Emerg Nurs. 1983;9:3141-144.*
19. *Berger ME, Jones OW, Ricks RC, Garrett S. Decontaminating the nasal passages. Operational Radiation Safety, Supplement to Health Physics. 2003;84:S80-S82.*
20. *Gerber GB, Thomas RG (eds). Guidebook for the treatment of accidental internal radionuclide contamination of workers. Radiation Protection Dosimetry. 1992:41:5-49.*

21. *US Department of Health and Human Services, Food and Drug Administration. Guidance for Industry, Internal Radioactive Contamination – Development of Decorporation Agents. March 2006. Available at <http://www.fda.gov/cder/guidance/6983fnl.htm> Accessed: May 22, 2006.*
22. *US Department of Health and Human Services, Food and Drug Administration. Guidance: potassium iodide as a thyroid blocking agent in radiation emergencies. December 2001. Available at: <http://www.fda.gov/cder/guidance/4825fnl.htm> Accessed: May 22, 2006.*
23. *National Research Council of the National Academies. Distribution and administration of potassium iodide in the event of a nuclear incident. Washington, DC: The National Academies Press; 2004.*
24. *Waselenko JK, MacVittie TJ, Blakely WF, et al. Medical management of the acute radiation syndrome: recommendations of the strategic national stockpile radiation working group. Ann Int Med. 2004; 140: 1037-1051.*
25. *Hughes WT, Armstrong D, Bodey GP, et al. 1997 Guidelines for the use of antimicrobial agents in neutropenic patients with unexplained fever. Clin Infectious Diseases. 1997;25:551-73.*
26. *International Commission on Radiological Protection Publication 60. 1990 Recommendations of the International Commission on Radiological Protection. New York: Elsevier Science; 1990.*
27. *United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR). Hereditary effects of radiation, report to the General Assembly, with scientific annexes No. E.01.1X.2. New York: United Nations; 2001, 1-2.*
28. *Mossman KL, Hill LT. Radiation risks in pregnancy. Obstet Gynecol. 1982;60:237-42.*
29. *Doll R, Wakeford F. Risk of childhood cancer from fetal irradiation. Br J Radiol. 1997;70:130-139.*
30. *Hall EJ. Radiobiology for the Radiologist, 5th edition. Philadelphia: Lippincott Williams & Wilkins; 2000, 189-191.*

31. *Mettler FA, Upton AC. Medical effects of ionizing radiation, 2nd edition. Philadelphia: WB Saunders, 1995, 329.*
32. *Annals of the International Commission on Radiological Protection, Publication 84. Pregnancy and medical radiation. Tarrytown: Elsevier Science Inc.; 2000.*